

Transport and Noise of Entangled Electrons

Eugene V. Sukhorukov, Daniel Loss, and Guido Burkard

Department of Physics and Astronomy, University of Basel,
Klingelbergstrasse 82, CH-4056 Basel, Switzerland

Abstract. We consider a scattering set-up with an entangler and beam splitter where the current noise exhibits bunching behavior for electronic singlet states and antibunching behavior for triplet states. We show that the entanglement of two electrons in the double-dot can be detected in mesoscopic transport measurements. In the cotunneling regime the singlet and triplet states lead to phase-coherent current contributions of opposite signs and to Aharonov-Bohm and Berry phase oscillations in response to magnetic fields. We analyze the Fermi liquid effects in the transport of entangled electrons.

1 Introduction

The availability of pairwise entangled qubits – Einstein-Podolsky-Rosen (EPR) pairs [1] – is a necessary prerequisite in quantum communication [2]. The prime example of an EPR pair considered here is the singlet/triplet state formed by two electron spins [3], [4]. Its main feature is its non-locality: If we separate the two electrons from each other in real space, their total spin state can still remain entangled. Such non-locality gives rise to striking phenomena such as violations of Bell inequalities and quantum teleportation and has been investigated for photons [5], [6], but not yet for *massive* particles such as electrons, let alone in a solid state environment. In this work we discuss specific properties of transport and noise of entangled electrons as a result of two-particle coherence and nonlocality.

In Sect. (2) we propose and analyze an experimental set-up (see Fig. 1a) by which the entanglement of mobile electrons can be detected in noise measurements with a beam splitter [7]. The entangler is assumed to be a device by which we can generate entangled electron states, a specific realization being the double-dot system [3]. The presence of a beam splitter ensures that the electrons leaving the entangler have a finite amplitude to be interchanged. Thus we can expect that the current-current correlations (noise) measured in leads 3 and/or 4 are sensitive to the symmetry of the *orbital part* of the wave function [8]. Since the spin singlet of two electrons is uniquely associated with a symmetric orbital wave-function, and the triplet with an antisymmetric one we have a means to distinguish singlets from triplets through a bunching or antibunching signature.

[†] To appear in the Proceedings of the XVI Sitges Conference, Statistical and Dynamical Aspects of Mesoscopic Systems, Lecture Notes in Physics, Springer.




Fig. 1. a) The setup for measuring noise correlations of entangled states. Uncorrelated electrons are fed into the entangler through the Fermi leads $1'$ and $2'$. The entangler is a device that produces pairs of entangled electrons and injects one of the electrons into lead 1 and the other into lead 2. The entanglement can then be detected by performing an interference experiment using a beam splitter. b) Double-dot (DD) system containing two electrons and being weakly coupled to metallic leads 1,...,4, each of which being at the chemical potential $\mu_1, \dots, \mu_4$. The tunneling amplitudes between dots and leads are denoted by Γ , $\tilde{\Gamma}$. The tunneling (t) between the dots results in a singlet (triplet) ground state. The closed tunneling path between dots and leads 1 and 2 encloses the area A .

It is well-known [10] that bosons (fermions) show bunching (antibunching) behavior [11]. Antibunching is so far considered for electrons in the normal state both in theory [12], [13] and in experiments [14]. However, this classical effect is independent of phase coherence [15] and should be carefully distinguished from the two-particle phase-coherent effect which we propose here.

The scheme we propose in Sect. (3) [16] consists of two coupled quantum dots (DD) which themselves are weakly coupled in parallel to two leads 1 and 2 (see Fig. 1b). This results in a closed loop, and applying a magnetic field, an Aharonov-Bohm (AB) phase φ will be accumulated by an electron traversing the DD. In the Coulomb blockade (CB) regime we find that due to cotunneling [17] the current depends on the state of the DD: the AB oscillations for singlet and triplets have opposite sign. The amplitude of the AB oscillations provides a measure of the phase coherence of the entangled state, while the period – via the enclosed area A – provides a measure of the non-locality of the EPR pairs. The triplets themselves can be further distinguished by applying a directionally inhomogeneous magnetic field which adds a Berry phase [18] leading to beating.

Finally, in Sect. (4) we would like to address the following question [9]: Is

it possible to use mobile electrons, prepared in an entangled spin state, for the purpose of quantum communication? Without spin-dependent interaction we know that the total spin must be conserved even if the two electrons strongly interact with the other electrons in the mesoscopic environment (and among themselves) via Coulomb interaction. It is thus not unreasonable to expect that we still find some spin correlations between initial and final states. But how much is it? And why and how do we lose some of the correlations?

2 Noise of Entangled Electrons: Beam Splitter Set-up

Below, we extend the standard scattering matrix approach [12] to a situation with entanglement. We start by writing the operator for the current carried by electrons with spin σ in lead α of a multiterminal conductor as

$$I_{\alpha\sigma}(t) = \frac{e}{h\nu} \sum_{\epsilon, \epsilon'} \sum_{\beta\gamma} a_{\beta\sigma}^\dagger(\epsilon) a_{\gamma\sigma}(\epsilon') A_{\beta\gamma}^\alpha \exp[i(\epsilon - \epsilon')t/\hbar] , \quad (1)$$

$$A_{\beta\gamma}^\alpha = \delta_{\alpha\beta} \delta_{\alpha\gamma} - s_{\alpha\beta}^* s_{\alpha\gamma} . \quad (2)$$

where $a_{\alpha\sigma}^\dagger(\epsilon)$ creates an incoming electron in lead α with spin σ and energy ϵ , and we assume that the scattering matrix $s_{\alpha\beta}$ is spin- and energy-independent. Note that since we are dealing with discrete energy states here, we normalize the operators $a_\alpha(\epsilon)$ such that $[a_{\alpha\sigma}(\epsilon), a_{\beta\sigma'}^\dagger(\epsilon')] = \delta_{\sigma\sigma'} \delta_{\alpha\beta} \delta_{\epsilon, \epsilon'}/\nu$, where the Kronecker symbol $\delta_{\epsilon, \epsilon'}$ equals 1 if $\epsilon = \epsilon'$ and 0 otherwise. Therefore we also have included the factor $1/\nu$ in the definition of the current, where ν stands for the density of states in the leads. We will also assume that each lead consists of only a single quantum channel; the generalization to leads with several channels is straightforward but is not needed here.

We restrict ourselves here to unpolarized currents, $I_\alpha = \sum_\sigma I_{\alpha\sigma}$. The spectral density current fluctuations (noise) $\delta I_\alpha = I_\alpha - \langle I_\alpha \rangle$ between the leads α and β are defined as

$$S_{\alpha\beta}(\omega) = \lim_{T \rightarrow \infty} \frac{h\nu}{T} \int_0^T dt \exp(i\omega t) \langle \Psi | \delta I_\alpha(t) \delta I_\beta(0) | \Psi \rangle , \quad (3)$$

where $|\Psi\rangle$ is the quantum state of the system to be specified next¹. We will now investigate the noise for scattering with the entangled incident state

$$|\pm\rangle = \frac{1}{\sqrt{2}} \left(a_{2\downarrow}^\dagger(\epsilon_2) a_{1\uparrow}^\dagger(\epsilon_1) \pm a_{2\uparrow}^\dagger(\epsilon_2) a_{1\downarrow}^\dagger(\epsilon_1) \right) |0\rangle . \quad (4)$$

The state $|-\rangle$ is the spin singlet, $|S\rangle$, while $|+\rangle$ denotes the $S_z = 0$ triplet $|T_0\rangle$ ². Substituting $|\pm\rangle$ defined in (4) for $|\Psi\rangle$ and using the fact that the unpolarized currents are invariant when all spins are reversed, the expectation value

¹ Note that since $|\Psi\rangle$ in general does not describe a Fermi liquid state, it is not possible to apply Wick's theorem.

² All three triplets, $|T_0\rangle$, $|\uparrow\uparrow\rangle$, $|\downarrow\downarrow\rangle$, give the same result.

$\langle \pm | \delta I_\alpha \delta I_\beta | \pm \rangle$ can be expressed as the sum of a direct and an exchange term,

$$\langle \pm | \delta I_\alpha \delta I_\beta | \pm \rangle = \langle \uparrow \downarrow | \delta I_\alpha \delta I_\beta | \uparrow \downarrow \rangle \pm \langle \uparrow \downarrow | \delta I_\alpha \delta I_\beta | \downarrow \uparrow \rangle, \quad (5)$$

where the upper (lower) sign of the exchange term refers to triplet (singlet). Direct calculation of (5) gives the following result for the zero-frequency ($\omega = 0$) correlations,

$$S_{\alpha\beta} = \frac{e^2}{h\nu} \left[\sum_{\gamma\delta}' A_{\gamma\delta}^\alpha A_{\delta\gamma}^\beta \mp \delta_{\epsilon_1, \epsilon_2} (A_{12}^\alpha A_{21}^\beta + A_{21}^\alpha A_{12}^\beta) \right], \quad (6)$$

where $\sum_{\gamma\delta}'$ denotes the sum over $\gamma = 1, 2$ and all $\delta \neq \gamma$.

We apply formula (6) now to the set-up shown in Fig. 1a involving four leads, described by the single-particle scattering matrix elements, $s_{31} = s_{42} = r$, and $s_{41} = s_{32} = t$, where r and t denote the reflection and transmission amplitudes at the beam splitter, respectively. We assume that there is no backscattering, $s_{12} = s_{34} = s_{\alpha\alpha} = 0$. The unitarity of the s-matrix implies $|r|^2 + |t|^2 = 1$, and $\text{Re}[r^*t] = 0$. Using (2) and (6), we obtain the final result for the noise correlations for the incident states $|\pm\rangle$,

$$\begin{aligned} S_{33} &= S_{44} = -S_{34} = eF |\langle I \rangle|, \\ F &= 2eT(1 - T) (1 \mp \delta_{\epsilon_1, \epsilon_2}), \end{aligned} \quad (7)$$

where $|\langle I \rangle| = e/h\nu$ is the average current in all leads, $T = |t|^2$ is the probability for transmission through the beam splitter, and F is the Fano factor. Note that the total current $\delta I_3 + \delta I_4$ does not fluctuate, i.e. $S_{33} + S_{44} + 2S_{34} = 0$, since we have excluded backscattering.

Above results (7) imply that if two electrons with the same energies, $\epsilon_1 = \epsilon_2$, in the singlet state $|S\rangle = |-\rangle$ are injected into the leads 1 and 2, then the zero frequency noise is *enhanced* by a factor of two, $F = 4eT(1 - T)$, compared to the shot noise of uncorrelated particles, $F = 2eT(1 - T)$. This enhancement of noise is due to *bunching* of electrons in the outgoing leads, caused by the symmetric orbital wavefunction of the spin singlet $|S\rangle$. On the other hand, all three triplets $|+\rangle$ exhibit an *antibunching* effect, leading to a complete suppression of the zero-frequency noise, $S_{\alpha\alpha} = 0$. The noise enhancement for the singlet $|S\rangle$ is a unique signature for entanglement (there exists no unentangled state with the same symmetry), therefore entanglement can be observed by measuring the noise power of a mesoscopic conductor as shown in Fig. 1a.³

3 Probing Entanglement of Electrons in a Double-Dot

The DD system (see Fig. 1b) contains 4 metallic leads which are in equilibrium with associated reservoirs kept at the chemical potentials μ_i , $i = 1, \dots, 4$. The

³ These results remain valid in the presence of a Fermi sea.

leads are weakly coupled to the dots with tunneling amplitudes Γ and $\tilde{\Gamma}$, and the leads 1, 2 are coupled to *both* dots and play the role of probes where the currents I_i are measured. The leads 3 and 4 are feeding electrodes to manipulate the electron filling in the dots. The quantum dots contain one (excess) electron each, and are coupled to each other by the tunneling amplitude t , which leads to a level splitting [3], [4] $J = E_t - E_s \sim 4t^2/U$ in the DD, with U being the single-dot Coulomb repulsion energy, and $E_{s/t}$ are the singlet/triplet energies. We recall that for two electrons in the DD (and for weak magnetic fields) the ground state is given by a spin singlet. For convenience we count the chemical potentials μ_i from E_s . The coupling $\tilde{\Gamma}$ to the feeding leads can be switched off while probing the DD with a current. Here we assume that $\tilde{\Gamma} = 0$.

Using a standard tunneling Hamiltonian approach [19], we write $H = H_0 + V$, where the first term in $H_0 = H_D + H_1 + H_2$ describes the DD and $H_{1,2}$ the leads (assumed to be Fermi liquids). The tunneling between leads and dots is described by the perturbation $V = V_1 + V_2$, where

$$V_n = \Gamma \sum_s [D_{n,s}^\dagger c_{n,s} + c_{n,s}^\dagger D_{n,s}], \quad D_{n,s} = e^{\pm i\varphi/4} d_{1,s} + e^{\mp i\varphi/4} d_{2,s}, \quad (8)$$

and where the operators $c_{n,s}$ and $d_{n,s}$, $n = 1, 2$, annihilate electrons with spin s in the n th lead and in the n th dot, resp. The Peierls phase φ in the hopping amplitude accounts for an AB or Berry phase (see below) in the presence of a magnetic field. The upper sign belongs to lead 1 and the lower to lead 2. Finally, we assume that spin is conserved in the tunneling process. For the outgoing currents we have $I_n = ie\Gamma \sum_s [D_{n,s}^\dagger c_{n,s} - c_{n,s}^\dagger D_{n,s}]$. The observable of interest is the average current through the DD system, $I = \langle I_2 \rangle$.

From now on we concentrate on the CB regime where we can neglect double (or higher) occupancy in each dot for all transitions including virtual ones, i.e. we require $\mu_{1,2} < U$. Further we assume that $\mu_{1,2} > J, k_B T$ to avoid resonances which might change the DD state. The lead-dot coupling Γ is assumed to be weak so that the state of the DD is not perturbed; this will allow us to retain only the first non-vanishing contribution in Γ to I . Formally, we require $J > 2\pi\nu_t \Gamma^2$, where ν_t is the tunneling density of states of the leads. In analogy to the single-dot case [17] we refer to above CB regime as cotunneling regime.

Continuing with our derivation of I , we note that the average $\langle \dots \rangle \equiv \text{Tr} \rho \{ \dots \}$ is taken with respect to the equilibrium state of the *entire* system set up in the distant past before V is switched on [19]. Then, in the interaction picture, the current is given by

$$I = \langle U^\dagger I_2(t) U \rangle, \quad U = T \exp \left[-i \int_{-\infty}^t dt' V(t') \right]. \quad (9)$$

The leading contribution in Γ to the cotunneling current involves the tunneling of one electron from the DD to, say, lead 2 and of a second electron from lead 1 to the DD (see Fig. 1b). This contribution is of order $V_2 V_1^2$, and thus $I \propto \Gamma^4$, as

is typical for cotunneling [17]. Taking the trace over Fermi leads, we arrive then at the following compact expression for the cotunneling current

$$I = \frac{1}{2} e \pi \nu_t^2 \Gamma^4 \sum_{i,f,s,s'} \rho_i |\langle i | D_{2,s'}^\dagger D_{1,s} | f \rangle|^2 \frac{\Delta_{i,f} \theta(\Delta_{i,f})}{\mu_1 \mu_2} ,$$

$$\Delta_{i,f} = \mu_1 - \mu_2 + E_i - E_f . \quad (10)$$

This equation shows that in the cotunneling regime the initial state $|i\rangle$ (with weight ρ_i) of the DD is changed into a final state $|f\rangle$ by the traversing electron. However, due to the weak coupling Γ , the DD will have returned to its equilibrium state before the next electron passes through it.

For small bias, $|\mu_1 - \mu_2| < J$, only elastic cotunneling is allowed, i.e. $E_i = E_f$. However, this regime is not of interest here since singlet and triplet contributions turn out to be identical and thus indistinguishable. We thus focus on the opposite regime, $|\mu_1 - \mu_2| > J$, where inelastic cotunneling⁴ occurs with singlet and triplet contributions being different. In this regime we can neglect the dynamics generated by J compared to the one generated by the bias (“slow spins”), and drop the energies E_i and E_f in (10). Finally, using $1 = \sum_f |f\rangle\langle f|$ we obtain

$$I = e \pi \nu_t^2 \Gamma^4 C(\varphi) \frac{\mu_1 - \mu_2}{\mu_1 \mu_2} , \quad (11)$$

$$C(\varphi) = \sum_{s,s'} \left[\langle d_{1s'}^\dagger d_{1s} d_{1s}^\dagger d_{1s'} \rangle + \cos \varphi \langle d_{1s'}^\dagger d_{1s} d_{2s}^\dagger d_{2s'} \rangle \right] . \quad (12)$$

For the purpose of our analysis we assume that the DD is in its ground state. Equation (12) shows that the cotunneling current depends on the properties of the ground state of the DD through the coherence factor $C(\varphi)$ given in (12). The first term in C is the contribution from the topologically trivial tunneling path which runs from lead 1 through, say, dot 1 to lead 2 and back. The second term (phase-coherent part) in C is the ground state amplitude of the exchange of electron 1 with electron 2 via the leads 1 and 2 such that a closed loop is formed enclosing an area A (see Fig. 1b). Thus, in the presence of a magnetic field B , an AB phase factor $\varphi = ABe/h$ is acquired.

Next, we evaluate $C(\varphi)$ explicitly in the singlet-triplet basis. Note that only the singlet $|S\rangle$ and the triplet $|T_0\rangle$ (see (4)) are entangled EPR pairs while the remaining triplets $|T_+\rangle = |\uparrow\uparrow\rangle$, and $|T_-\rangle = |\downarrow\downarrow\rangle$ are not (they factorize). Assuming that the DD is in one of these states we obtain the important result

$$C(\varphi) = \begin{cases} 2 - \cos \varphi , & \text{for singlet} \\ 2 + \cos \varphi , & \text{for all triplets} . \end{cases} \quad (13)$$

Thus, we see that the singlet and the triplets contribute with *opposite sign to the phase-coherent part of the current*. One has to distinguish, however, carefully

⁴ Note that the AB effect is not suppressed by this inelastic cotunneling, since the *entire* cotunneling process involving also leads is elastic: the initial and final states of the *entire* system have the same energy.

the entangled from the non-entangled states. The phase-coherent part of the entangled states is a genuine *two-particle* effect, while the one of the product states cannot be distinguished from a phase-coherent *single-particle* effect. Indeed, this follows from the observation that the phase-coherent part in C factorizes for the product states $T_{\pm}$ while it does not so for S, T_0 . Also, for states such as $|\uparrow\downarrow\rangle$ the coherent part of C vanishes, showing that two different (and fixed) spin states cannot lead to a phase-coherent contribution since we *know* which electron goes which part of the loop. Finally we note that due to the AB phase the role of the singlet and triplets can be interchanged which is to say that we can continually transmute the statistics of the entangled pairs S, T_0 from fermionic to bosonic (like in anyons): the symmetric orbital wave function of the singlet S goes into an antisymmetric one at half a flux quantum, and vice versa for the triplet T_0 .

We would like to stress that the amplitude of the AB oscillations is a direct measure of the phase coherence of the entanglement, while the period via the enclosed area $A = h/eB_0$ gives a direct measure of the non-locality of the EPR pairs, with B_0 being the field at which $\varphi = 1$. The triplets themselves can be further distinguished by applying a directionally inhomogeneous magnetic field (around the loop) producing a Berry phase Φ^B [18], which is positive (negative) for the triplet $m = 1(-1)$, while it vanishes for the EPR pairs S, T_0 . Thus, we will eventually see beating in the AB oscillations due to the positive (negative) shift of the AB phase Φ by the Berry phase, $\varphi = \Phi \pm \Phi^B$.

4 Transport of Entangled Electrons

We consider the general scenario of the transport of entangled electrons in a mesoscopic system [9]. In a first step we inject entangled electrons into the leads and create the state $|\mathbf{12}\rangle \equiv |\pm\rangle$ (see (4)) on the top of the Fermi sea (as discussed e.g. in Sect. (2), see Fig. 1a). In a second step, we perform a quantum measurement of the state. As a measure of correlations we consider transition amplitudes between an initial and a final state. We begin with the simplest case given by the wave function overlap of $|\mathbf{12}\rangle$ with $|\mathbf{34}\rangle$,

$$\langle \mathbf{12} | \mathbf{34} \rangle = \delta_{\mathbf{13}} \delta_{\mathbf{24}} \mp \delta_{\mathbf{14}} \delta_{\mathbf{23}} \quad , \quad (14)$$

where the upper (lower) sign refers to triplet (singlet). If the quantum numbers coincide, $\mathbf{1} = \mathbf{3}$, and $\mathbf{2} = \mathbf{4}$, the overlap assumes its maximum value 1, reflecting maximum correlation between the two states.

Next we generalize this concept to leads which contain many interacting electrons besides the two entangled electrons. We use a similar overlap as a measure of how much weight remains in the final state $|\mathbf{34}, t\rangle$ when we start from some given initial state $|\mathbf{12}\rangle$. The overlap (14) now becomes a triplet-triplet or singlet-singlet correlation function

$$G^{t/s}(\mathbf{12}, \mathbf{34}; t) = -G(\mathbf{1}, t) G(\mathbf{2}, t) [\delta_{\mathbf{13}} \delta_{\mathbf{24}} \mp \delta_{\mathbf{14}} \delta_{\mathbf{23}}] \quad , \quad (15)$$

where we have assumed that there is no interactions between lead 1 and 2. Thus the problem is reduced to the evaluation of (time-ordered) single-particle Green's functions $G(\mathbf{1}, t)$, $G(\mathbf{2}, t)$ pertaining to lead 1 and 2, resp. (these leads are still interacting many-body systems though).

For the special case $t = 0$, and no interactions, we have $G = -i$, and thus $G^{t/s}$ reduces to the rhs of (14). For the general case, we evaluate G close to the Fermi surface and get the standard result [20]

$$G(\epsilon, t) \approx -iz_\epsilon \Theta(\epsilon - \epsilon_F) e^{-i\epsilon t - \Gamma_\epsilon t} , \quad (16)$$

where ϵ is the quasiparticle energy, ϵ_F is the Fermi energy, and $1/\Gamma_\epsilon$ is the quasiparticle lifetime. In a 2DEG, $\Gamma_\epsilon \propto (\epsilon - \epsilon_F)^2 \log(\epsilon - \epsilon_F)$ [21] within the random phase approximation (RPA). Thus, the lifetime becomes infinite when the energy of the added electron approaches ϵ_F .

Now, we come to the most important quantity in the present context, the *quasiparticle weight*, $z_F = z_{\epsilon_F}$, evaluated at the Fermi surface; it is defined by

$$z_F = \left[1 - \frac{\partial}{\partial \omega} \text{Re} \Sigma(\epsilon_F, \omega = 0) \right]^{-1} , \quad (17)$$

where $\Sigma(\epsilon, \omega)$ is the irreducible self-energy occurring in the Dyson equation. The quasiparticle weight, $0 \leq z_\epsilon \leq 1$, describes the weight of the bare electron in the quasiparticle state ϵ , i.e. when we add an electron with energy $\epsilon \geq \epsilon_F$ to the system, some weight (given by $1 - z_\epsilon$) of the original state ϵ will be distributed among all the electrons due to the Coulomb interaction.

Restricting ourselves now to energies close to the Fermi surface we have

$$G^{t/s}(\mathbf{12}, \mathbf{34}; t) = z_F^2 [\delta_{13} \delta_{24} \mp \delta_{14} \delta_{23}] , \quad (18)$$

for all times satisfying $0 < t \lesssim 1/\Gamma_\epsilon$. Thus we see that it is the quasiparticle weight squared, z_F^2 , which is the measure of our spin correlation function $G^{t/s}$ we were looking for. It is thus interesting to evaluate z_F explicitly. This is indeed possible, again within RPA, and we find after careful calculation [22]

$$z_F = 1 - r_s \left(\frac{1}{2} + \frac{1}{\pi} \right) , \quad (19)$$

in leading order of the interaction parameter $r_s = 1/q_F a_B$, where $a_B = \epsilon_0 \hbar^2 / m e^2$ is the Bohr radius. In particular, in a GaAs 2DEG we have $a_B = 10.3$ nm, and $r_s = 0.614$, and thus we obtain from (19) the value $z_F = 0.665$. We note that a more accurate numerical evaluation of the exact RPA self-energy yields⁵ $z_F = 0.691$ [22]. Thus, we see that the spin correlation is reduced by a factor of about two as soon as we inject the two electrons into separate leads consisting of *interacting* Fermi liquids in their ground state.

⁵ For 3D metallic leads with, say, $r_s = 2$ (e.g. $r_s^{\text{Cu}} = 2.67$) the loss of correlation is somewhat less strong, since then the quasiparticle weight becomes $z_F = 0.77$ [19].

5 Acknowledgements

This work has been supported by the Swiss National Science Foundation.

References

- [1] A. Einstein, B. Podolsky, N. Rosen Phys. Rev. **47**, 777 (1935).
- [2] C. H. Bennett and G. Brassard, in Proceedings of the IEEE International Conference on Computers, Systems and Signal Processing, Bangalore, India (IEEE, New York, 1984), p. 175.
- [3] D. Loss, D. P. DiVincenzo, Phys. Rev. A **57**, 120 (1998).
- [4] G. Burkard, D. Loss, D. P. DiVincenzo, Phys. Rev. B **59** 2070 (1999).
- [5] A. Aspect, J. Dalibard, and G. Roger, Phys. Rev. Lett. **49**, 1804 (1982).
- [6] A. Zeilinger, in *Physics World*, March 1998.
- [7] G. Burkard, D. Loss and E. V. Sukhorukov, preprint (cond-mat/9906071).
- [8] R. P. Feynman, R. B. Leighton, and M. Sands, *The Feynman lectures on physics*, Vol. 3 (Addison-Wesley, 1965).
- [9] D. P. DiVincenzo, D. Loss, preprint (cond-mat/9901137).
- [10] R. Loudon, in *Coherence and Quantum Optics VI*, eds. J. H. Eberly et al. (Plenum, New York, 1990).
- [11] R. Hanbury Brown and R. Q. Twiss, Nature (London) **177**, 27 (1956).
- [12] M. Büttiker, Phys. Rev. Lett. **65**, 2901 (1990); Phys. Rev. B **46**, 12485 (1992).
- [13] T. Martin, R. Landauer, Phys. Rev. B **45**, 1742 (1992).
- [14] R. C. Liu et al., Nature **391**, 263 (1998); M. Henny et al., Science **284**, 296 (1999); W. D. Oliver et al., Science **284**, 299 (1999).
- [15] E. V. Sukhorukov and D. Loss, Phys. Rev. B **59**, 13054 (1999).
- [16] D. Loss and E. V. Sukhorukov, preprint (cond-mat/9907129).
- [17] D. V. Averin, Yu. V. Nazarov, in *Single Charge Tunneling*, eds. H. Grabert and M.H. Devoret, NATO ASI Series B: Physics Vol. 294, Plenum Press, New York, 1992; J. König, H. Schoeller, and G. Schön, Phys. Rev. Lett. **78**, 4482 (1997).
- [18] D. Loss, P. Goldbart, Phys. Rev. B **45**, 13544 (1992).
- [19] G. D. Mahan, *Many Particle Physics*, 2nd Ed. (Plenum, New York, 1993).
- [20] A.L. Fetter and J.D. Walecka, *Quantum Theory of Many-Particle Systems* (New York, McGraw-Hill, 1971).
- [21] G. F. Giuliani and J. J. Quinn, Phys. Rev. B **26**, 4421 (1982).
- [22] D. Loss, G. Burkard, and E. Sukhorukov, *Quantum communication with electrons*, unpublished.